Hamiltonian Mechanics

Ivar Ekeland[1] and Roger Temam[2]

[1] Princeton University, Princeton NJ 08544, USA
[2] Université de Paris-Sud, Laboratoire d'Analyse Numérique, Bâtiment 425, F-91405 Orsay Cedex, France

Abstract. The abstract should summarize the contents of the paper using at least 70 and at most 150 words. It will be set in 9-point font size and be inset 1.0 cm from the right and left margins. There will be two blank lines before and after the Abstract. ...

1 Fixed-Period Problems: The Sublinear Case

With this chapter, the preliminaries are over, and we begin the search for periodic solutions to Hamiltonian systems. All this will be done in the convex case; that is, we shall study the boundary-value problem

$$\begin{aligned} \dot{x} &= JH'(t,x) \\ x(0) &= x(T) \end{aligned}$$

with $H(t,\cdot)$ a convex function of x, going to $+\infty$ when $\|x\| \to \infty$.

1.1 Autonomous Systems

In this section, we will consider the case when the Hamiltonian $H(x)$ is autonomous. For the sake of simplicity, we shall also assume that it is C^1.

We shall first consider the question of nontriviality, within the general framework of (A_∞, B_∞)-subquadratic Hamiltonians. In the second subsection, we shall look into the special case when H is $(0, b_\infty)$-subquadratic, and we shall try to derive additional information.

The General Case: Nontriviality. We assume that H is (A_∞, B_∞)-subquadratic at infinity, for some constant symmetric matrices A_∞ and B_∞, with $B_\infty - A_\infty$ positive definite. Set:

$$\gamma := \text{smallest eigenvalue of } \; B_\infty - A_\infty \tag{1}$$

$$\lambda := \text{largest negative eigenvalue of } \; J\frac{d}{dt} + A_\infty \;. \tag{2}$$

Theorem 21 tells us that if $\lambda + \gamma < 0$, the boundary-value problem:

$$\begin{aligned} \dot{x} &= JH'(x) \\ x(0) &= x(T) \end{aligned} \tag{3}$$